

CEREALS

PRIORITY OF ADMINISTRATION ACCORDING TO STAGES OF PLANT DEVELOPMENT ■ required ■ important ■ optional

0,1 kg/t	To the leaf together with pesticides: 1-4 times 0,06-0,08 kg/ha							
	7	21	25	29	30	51-59	61-69	71-75
Seed treatment	seedlings		tillering		Stem elongation	Ear formation	flowering	Milk stage

RE:GREENER is effective from first half of vegetation until the flowering, especially in rosette and stooling stages. It is useful to add 0,06 kg/ha of RE:GREENER for every treatment and PPP to decrease the pesticide load for plants.



CANOLA

0,1 kg/t	To the leaf: 2-4 treatments 0,06-0,2 kg/ha						
	0-10	18-19	21-22	31-39	51-57	58-69	70-75
Seed treatment	seedlings	3-4 leaves	rosette	stooling	budding	flowering	Milk stage

RE:GREENER is effective from first half of vegetation until the flowering, especially in rosette and stooling stages. It is useful to add 0,06 kg/ha of RE:GREENER for every treatment and PPP to decrease the pesticide load for plants.



RICE

0,1 kg/t	To the leaf: 1-2 times 0,15-0,2 kg/ha							
	7	21	25	29	30	51-59	61-69	71-75
Seed treatment	seedlings		tillering		Stem elongation	Ear formation	flowering	Milk stage

Treatment of seeds is as important agronomic practice, ensuring the increase development of root system. When treating with carbamide or KAS to the leaf – for increased effect add RE:GREENER in tank mixture (treatment effect increasing up to 50%). Increase of crop-producing depending from the scheme and agricultural background from 5 to 25 cwt/ha.



MAIZE

PRIORITY OF ADMINISTRATION ACCORDING TO STAGES OF PLANT DEVELOPMENT ■ required ■ important ■ optional

0,1 kg/t	To the leaf together with pesticides: 1-2 times 0,1-0,2 kg/ha							
	7	21	25	29	30	51-59	61-69	71-75
Seed treatment	seedlings	3-5 leaves		6-8 leaves		tasseling	flowering	Milk stage


The most important – treatment of seeds and treatment at 3-7 leaf stage together with herbicides and fertilizers. It is essential to add 0,05 kg/ha of RE:GREENER at every spraying with PPP to decrease the chemical load for the plants. Increase of crop-producing depending from the scheme and agricultural background from 7 to 10 cwt/ha.



PULSES soy, peas, pinto

0,1 kg/t	To the leaf: 2-3 times 0,06-0,1 kg/ha					
	0-10	12-15	50-53	54-57	58-69	70-79
Seed treatment	seedlings	1-3 leaves	tillering	budding	flowering	Milk stage

RE:GREENER is effective during the whole vegetation, for foliar application together with Aquarine and Magnesium Sulfate. Ideal in treatment of seeds with biological drugs (Risotorpine etc.) increasing the effectiveness of Nitrogen fixation. It is useful to add 0,05 kg/ha of RE:GREENER 2-3 times per vegetation, including in tank mixtures with PPP to decrease the chemical load for the plants. Increase of crop-producing from 3,5 cwt/ha.



IMPORTANT TO BE AWARE.

- It is recommended to add RE:GREENER together with plant protection products (PPP), foliar applications, complex water-soluble, nitrogen fertilizers for increase of effectiveness of plant protection products, taking of macro and micro nutrients by plants and decrease of consumption thereof.
- It is recommended to perform the preparation of spray material solution right before use, especially in case of using hard water.
- In case of feeding through drip system use 0,02% of RE:GREENER solution (once per 10-14 days, at least 3-4 times during the vegetation period).
- It is recommended to add RE:GREENER to decrease the stress from plants after use of pesticides, after frost, hail, etc. (dosage: 30-150 g/ha for 200-300 litres of water.)
- It is not recommended to mix RE:GREENER with agents, containing the high dosages of calcium or with tank mixtures with pH less than 4,5.