

SWEET PEPPER, EGGPLANT

PRIORITY OF ADMINISTRATION ACCORDING TO STAGES OF PLANT DEVELOPMENT ■ required ■ important ■ optional

0,05 kg/10 l of water	To the leaf: 2-7 times 0,1-0,2 kg/ha						
	Seed treatment	seedlings	1-2 leaves	2-4 leaves	3-5 leaves	budding	flowering

RE:GREENER is especially effective in root-stock watering (and at early stages in the direct seeding), when plants are being set to the fixed place-watering of seedbeds (0,02% mixture) and after the setting to the fixed place with soil dressing up to the fruit stage. In case of multiple outside-root treatment PPP it is recommended to add 0,05kg/ha of RE:GREENER at each treatment to decrease the chemical load on the stage. RE:GREENER shall be added to the drip watering at vegetation stage (0,02% mixture) every 7-12 days. May be used together with biological drugs (increase the effectiveness thereof).



ONION, GARLIC

0,05 kg/10 l of water 0,03 kg/t	To the leaf: 3-5 times 0,05-0,15 kg/ha					
	Seed treatment Treatment of planting material	seedlings	1-2 leaves	2-4 leaves	4-6 leaves	Increase of vegetative mass

In case of multiple outside-root treatment PPP it is recommended to add 0,05kg/ha of RE:GREENER at each treatment to decrease the chemical load on the stage. RE:GREENER shall be added to the drip watering at vegetation stage (0,02% mixture) every 7-12 days.



TOMATO

PRIORITY OF ADMINISTRATION ACCORDING TO STAGES OF PLANT DEVELOPMENT ■ required ■ important ■ optional

0,05 kg/10 l of water	To the leaf: 2-7 times 0,1-0,2 kg/ha						
	Seed treatment	seedlings	1-2 leaves	2-4 leaves	3-5 leaves	budding	flowering

RE:GREENER is especially effective in root-stock watering (and at early stages in the direct seeding), when plants are being set to the fixed place-watering of seedbeds (0,02% mixture) and after he setting to the fixed place with soil dressing up to the fruit formation stage. In case of multiple outside-root treatment PPP it is recommended to add 0,05kg/ha of RE:GREENER at each treatment to decrease the chemical load on the stage. RE:GREENER shall be added to the drip watering at vegetation stage (0,02% mixture) every 7-12 days. May be used together with biological drugs (increase the effectiveness thereof).




CUCUMBER, ZUCCHINI

PRIORITY OF ADMINISTRATION ACCORDING TO STAGES OF PLANT DEVELOPMENT ■ required ■ important ■ optional

0,05 kg/10 l of water	To the leaf: 2-7 times 0,1-0,2 kg/ha						
	Seed treatment	seedlings	1-2 leaves	2-4 leaves	6-8 leaves	budding	flowering

RE:GREENER is especially effective in root-stock watering (and at early stages in the direct seeding), when plants are being set to the fixed place-watering of seedbeds (0,02% mixture) and after he setting to the fixed place with soil dressing up to the fruit formation stage. In case of multiple outside-root treatment PPP it is recommended to add 0,05kg/ha of RE:GREENER at each treatment to decrease the chemical load on the stage. RE:GREENER shall be added to the drip watering at vegetation stage (0,02% mixture) every 7-12 days. May be used together with biological drugs (increase the effectiveness thereof).



POTATO

PRIORITY OF ADMINISTRATION ACCORDING TO STAGES OF PLANT DEVELOPMENT ■ required ■ important ■ optional

0,05 kg/t	To the leaf: 2-5 times 0,15-0,25 kg/ha				
		0-10	15-35	36-59	59-65
Seed treatment	seedlings	Stooling stage	budding	flowering	Tubers filling

In case of multiple outside-root treatment PPP it is recommended to add 0,05kg/ha of RE:GREENER at each treatment to decrease the chemical load on the stage. RE:GREENER shall be added to the drip watering at vegetation stage (0,02% mixture) every 7-12 days.



IMPORTANT TO BE AWARE.

1. It is recommended to add RE:GREENER together with plant protection products (PPP), foliar applications, complex water-soluble, nitrogen fertilizers for increase of effectiveness of plant protection products, taking of macro and micro nutrients by plants and decrease of consumption thereof.
2. It is recommended to perform the preparation of spray material solution right before use, especially in case of using hard water.
3. In case of feeding through drip system use 0,02% of RE:GREENER solution (once per 10-14 days, at least 3-4 times during the vegetation period).
4. It is recommended to add RE:GREENER to decrease the stress from plants after use of pesticides, after frost, hail, etc. (dosage: 30-150 g/ha for 200-300 litres of water.)
5. It is not recommended to mix RE:GREENER with agents, contending the high dosages of calcium or with tank mixtures with pH less than 4,5.